11413
Darunavir and Ritonavir Tablets 400mg/100mg
2070403

1. NAME OF THE MEDICINAL PRODUCT
Darunavir and Ritonavir Tablets 400mg/100mg

POM | **Schedule: S2** | **NS2** | **PP**
Prescription only medicine-List I

2. QUALITATIVE AND QUANTITATIVE COMPOSITION
Each Film coated tablets contains 433.640 mg of Darunavir Ethanolate equivalent to 400 mg of Darunavir and Ritonavir USP 50 mg.

For the full list of excipients, see section 6.1

3. PHARMACEUTICAL FORM
Film Coated Tablets

Yellow, capsule shaped, bevel edged, biconvex film coated tablets of 20 mm x 9 mm

side and 'DB' on the other side.

4. Clinical Particulars

4.1 Therapeutic Indications
Darunavir and Ritonavir Tablets 400mg/100mg is indicated in combination with other antiretroviral medicines for the treatment of human immunodeficiency virus type 1 (HIV-1) infection in adults and adolescents weighing at least 40 kg who:

- have not previously been treated with a protease inhibitor (protease inhibitor-naive patients); see section 4.2) or
- have previously been treated with a protease inhibitor (protease inhibitor-experienced) but do not harbour darunavir resistance associated mutations (see below) and who have plasma HIV-1 RNA < 100 000 copies/mL and CD4+ cell count ≥ 100 × 10⁶cells/L (see sections 4.4 and 5.1).

Darunavir and Ritonavir Tablets 400mg/100mg is used for protease inhibitor-experienced patients when HIV-1 genotype testing is available. Darunavir resistance associated mutations include mutations in codons V11, V32, L32F, I47V, I50V, I54L, T74P, L76V, I84V and L89V.

See section 4.4 for advice on darunavir/ritonavir treatment in protease inhibitor-experienced patients when genotype testing is not available.

Consideration should be given to official guidelines for treatment of HIV-1 infection (e.g. those of the WHO).

4.2 Posology and Method of Administration
Therapy should be initiated by a health care provider experienced in the management of HIV infection.

Posology

Patients weighing at least 40 kg
In PI-naive patients and in PI-experienced patients without darunavir-resistance associated mutations and who have plasma HIV-1 RNA < 100 000 copies/mL and CD4+ cell count ≥ 100 × 10⁶cells/L (see section 4.1), the recommended dose of Darunavir and Ritonavir Tablets 400mg/100mg is:

2 tablets taken once daily with food at around the same time each day.

Darunavir and Ritonavir Tablets 400mg/100mg is not suitable for treating all other ART-experienced patients or if HIV -1 genotype testing is not available.

Missed doses
If a dose of Darunavir and Ritonavir Tablets 400mg/100mg is missed within 12 hours of the time it is usually taken, patients should be instructed to take the dose of Darunavir and Ritonavir Tablets 400mg/100mg with food as soon as possible. If more than 12 hours have passed after the time it is usually taken, patients should not take the missed dose but the patient should resume the usual dosing schedule.

If a patient vomits within 4 hours of taking Darunavir and Ritonavir Tablets 400mg/100mg , the patient should take another dose with food as soon as possible. If the patient vomits more than 4 hours after taking the medicine, the patient does not need to take another dose.

Special populations

Elderly
Limited information is available in this population, and therefore, Darunavir and Ritonavir Tablets 400mg/100mg should be used with caution in this age group (see sections 4.4 and 5.2).

Hepatic impairment
Darunavir is metabolised by the hepatic system. No dose adjustment is recommended in patients with mild (Child-Pugh Class A) or moderate (Child-Pugh Class B) hepatic impairment; however, Darunavir and Ritonavir Tablets 400mg/100mg should be used with caution in these patients. No pharmacokinetic data are available in patients with severe hepatic impairment. Severe hepatic impairment could result in an increase of darunavir exposure and therefore Darunavir and Ritonavir Tablets 400mg/100mg must not be used in patients with severe hepatic impairment (Child-Pugh Class C) (see sections 4.3, 4.4 and 5.2).

Renal impairment
No dose adjustment is required for Darunavir and Ritonavir Tablets 400mg/100mg in patients with renal impairment (see sections 4.4 and 5.2).

Paediatric population
Darunavir and Ritonavir Tablets 400mg/50mg is not suitable for children weighing less than 40 kg. Other formulations of darunavir/ritonavir may be required.

Darunavir/ritonavir should not be used in children below 3 years of age or weighing less than 15 kg (see section 5.3).

Pregnancy and postpartum
No dose adjustment required for darunavir/ritonavir during pregnancy and postpartum. (see sections 4.6 and 5.2).

Method of administration
Patients should be instructed to take darunavir/ritonavir within 30 minutes after a meal. The type of food does not affect the exposure to darunavir (see section 5.2).

4.3 Contraindications
Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

Patients with severe (Child-Pugh Class C) hepatic impairment.

Concomitant treatment with medicines listed below because they can decrease plasma concentrations of darunavir and ritonavir, which could lead to loss of therapeutic effect and development of resistance (see sections 4.4 and 4.5). The following should not be used with Darunavir and Ritonavir Tablets 400mg/100mg:

- lopinavir/ritonavir,
- the strong CYP3A inducers rifampicin and herbal preparations containing St John's wort (*Hypericum perforatum*).

Darunavir boosted with ritonavir inhibits the elimination of active substances that are highly dependent on CYP3A for clearance. Therefore, concomitant treatment with medicines for which elevated plasma concentrations are associated with serious or life-threatening side effects is contraindicated (see also section 4.5). These active substances include:

- alfuzosin
- amiodarone, bepridil, dronedarone, encainide, flecainide, ivabradine, propafenone, quinidine, ranolazine
- astemizole, terfenadine
- avanafil, sildenafil when used for the treatment of pulmonary arterial hypertension, vardenafil
- cisapride
- clorazepate, diazepam, estazolam, flurazepam, oral midazolam and triazolam
- colchicine, lurasidone, pimozide, quetiapine, sertindole
- colchicine when used in patients with renal or hepatic impairment
- dabigatran, ticagrelor
- dapoxetine
- domperidone
- ergot derivatives (e.g. dihydroergotamine, ergometrine, ergotamine, methylergonovine)
- elbasvir/grazoprevir, ombitasvir/paritaprevir/ritonavir
- fusidic acid
- halofantrine, lumefantrine
- lomitapide, ivabradine and simvastatin
- naloxegol
- pimozide, sertindole
- pethidine, piroxicam, propoxyphene

4.4 Special warnings and precautions for use

Protease inhibitor-experienced patients
When genotype testing is not feasible in protease inhibitor-experienced patients, darunavir/ritonavir 600 mg/100 mg twice daily is recommended. Other formulations should be used. Darunavir and Ritonavir Tablets 400mg/50mg is not suitable.

Transmission
While effective viral suppression with antiretroviral therapy has substantially reduces the risk of sexual transmission, a residual risk may remain. Precautions to prevent transmission should be taken in accordance with relevant guidelines.

Regular assessment of virological response is advised. In the setting of lack or loss of virological response, resistance testing should be performed.

Antiretroviral therapy-experienced patients – once daily dosing
Darunavir/ritonavir at a dose of 800 mg/100 mg once daily in ART-experienced patients should not be used in patients with one or more darunavir resistance associated mutations (RAMs) or HIV-1 RNA > 100 000 copies/mL or CD4+ cell count < 100 x 10⁶cells/L (see section 4.2). Only combinations with optimised background regimen with two or more nucleotide (or nucleoside) reverse transcriptase inhibitors (NRTIs) should be studied in this population. Data are limited in patients with HIV-1 clades other than B (see section 5.1).

Elderly
As information is limited on the use of darunavir/ritonavir in patients aged 65 years and over, Darunavir and Ritonavir Tablets 400mg/50mg should be used with care in elderly patients, reflecting the greater frequency of decreased hepatic function and of concomitant disease or other therapy (see sections 4.2 and 5.2).

Severe skin reactions
Severe skin reactions, which may be accompanied with fever and elevated transaminases, have been reported. DRESS (drug rash with eosinophilia and systemic symptoms) or Stevens-Johnson syndrome has been reported rarely (< 0.1%), and during post-marketing experience toxic epidermal necrolysis and acute generalised exanthematous pustulosis have been reported. Symptoms can include severe rash or rash accompanied by fever, general malaise, fatigue, muscle or joint aches, blisters, oral lesions, conjunctivitis, hepatitis and eosinophilia (see also section 4.8). Darunavir and Ritonavir Tablets 400mg/50mg must be discontinued immediately if signs or symptoms of severe skin reactions develop.

Rash occurred more commonly in treatment-experienced patients receiving regimens containing darunavir/ritonavir + raltegravir compared to patients receiving darunavir/ritonavir + other antiretroviral agents without darunavir (see section 4.8).

Darunavir contains a sulphonamide moiety. Darunavir and Ritonavir Tablets 400mg/50mg should be used with caution in patients with sulphonamide allergy.

Hepatotoxicity
Drug-induced hepatitis (e.g. acute hepatitis, cytolytic hepatitis) has been reported in 0.5% of patients receiving combination antiretroviral therapy with darunavir /ritonavir. Patients with liver dysfunction, including chronic active hepatitis B or C, have an increased risk for liver function abnormalities including severe and potentially fatal hepatic adverse reactions. In case of concomitant antiviral therapy for hepatitis B or C, please refer to the relevant product information for these medicines.

Appropriate laboratory testing should be conducted before starting therapy with darunavir/ritonavir and patients should be monitored during treatment. Increased AST/ALT monitoring should be considered in patients with underlying chronic hepatitis, cirrhosis, or in patients who have pre-treatment elevations of transaminases, especially during the first several months of darunavir/ritonavir treatment. If there is evidence of new or worsening liver dysfunction (including clinically significant elevation of liver enzymes and symptoms such as fatigue, anorexia, nausea, jaundice, dark urine, liver tenderness, hepatomegaly) in patients using darunavir/ritonavir, interruption or discontinuation of treatment should be considered promptly.

Hepatic impairment
The safety and efficacy of darunavir have not been established in patients with severe liver disorders and therefore Darunavir and Ritonavir Tablets 400mg/50mg is therefore contraindicated in patients with severe hepatic impairment. Due to an increase in the unbound darunavir plasma concentrations, this medicine should be used with caution in patients with mild or moderate hepatic impairment (see sections 4.2, 4.3 and 5.2).

Renal impairment
No special precautions or dose adjustments for Darunavir and Ritonavir Tablets 400mg/100mg are required in patients with renal impairment. As darunavir and ritonavir are highly bound to plasma proteins, it is unlikely that they will be significantly removed by haemodialysis or peritoneal dialysis. Therefore, no special precautions or dose adjustments are required in these patients (see sections 4.3 and 5.2).

Patients with haemophilia
There have been reports of increased bleeding, including spontaneous skin haematomas and haemarthrosis in patients with haemophilia type A and B treated with protease inhibitors. In some patients additional factor VIII was given. In more than half of the reported cases, protease inhibitor treatment was not interrupted or reinstituted if treatment had been discontinued. A causal relationship has been suggested but the mechanism of action has not been elucidated. Patients with haemophilia should, therefore, be made aware of the possibility of increased bleeding.

Weight, blood lipids and glucose
Weight and levels of blood lipids and glucose may increase during antiretroviral therapy. Such changes may in part be linked to disease control and lifestyle. For lipids, there is some evidence of a treatment effect, while for weight gain there is no strong evidence relating this to any particular treatment. For monitoring blood lipids and glucose consult established HIV treatment guidelines. Lipid disorders should be managed as clinically appropriate.

Osteonecrosis
Cases of osteonecrosis have been reported particularly in patients with advanced HIV disease or long-term exposure to combination antiretroviral therapy. The aetiology is considered to be multifactorial (including corticosteroid use, alcohol consumption, severe immunosuppression, higher body mass index). Patients should be advised to seek medical advice if they have joint aches and pain, joint stiffness or difficulty in movement.

Opportunistic infections
Patients receiving darunavir or any other antiretroviral therapy may still develop opportunistic infections and other complications of HIV infection. Patients should remain under close clinical observation by health care providers experienced in the treatment of these associated HIV diseases.

Immune reconstitution inflammatory syndrome
When starting combination antiretroviral therapy (CART) in patients with severe immune deficiency an inflammatory reaction to asymptomatic or residual opportunistic pathogens may emerge and cause serious clinical conditions or aggravate symptoms. Typically such reactions occur within the first weeks or months of starting CART. Relevant examples are cytomegalovirus retinitis, generalised or focal mycobacterial infections and pneumonia caused by *Pneumocystis jirovecii* (formerly known as *Pneumocystis carinii*). Any inflammatory symptoms should be evaluated and treated when necessary. Also, reactivation of herpes simplex and herpes zoster has been observed in clinical studies with darunavir/ritonavir.

Autoimmune disorders (such as Graves' disease and autoimmune hepatitis) have also been reported in the setting of immune reactivation; however, the time to onset is more variable and these events can occur many months after starting treatment (see section 4.8).

Interactions with medicines
Etravirine: in combination with Darunavir and Ritonavir Tablets 400mg/100mg once daily may result in sub-optimal trough concentration of darunavir. If etravirine is to be used, darunavir/ritonavir should be dosed twice daily in combination with etravirine in treatment-experienced patients (see section 4.2).

Life-threatening and fatal drug interactions have been reported in patients treated with colchicine and strong inhibitors of CYP3A and P-glycoprotein (P-gp; see sections 4.3 and 4.5).

Darunavir binds predominantly to α1-acid glycoprotein. This protein binding is concentration-dependent indicative for saturation of binding. Therefore, there is a potential for protein displacement of medicines that are highly bound to α1-acid glycoprotein (see section 4.5).

4.5 Interactions
Interaction studies have only been performed in adults.

Darunavir and ritonavir are inhibitors of CYP3A, CYP2D6 and P-gp. Co-administration of darunavir/ritonavir with medicines primarily metabolised by CYP3A or CYP2D6 or transported by P-gp may increase systemic exposure to such medicines, which could increase or prolong their therapeutic effect and adverse reactions.

Coadministration of darunavir/ritonavir with drugs that have active metabolite(s) formed by CYP3A may result in reduced plasma concentrations of these active metabolite(s), potentially leading to loss of their therapeutic effect (see the interaction table below).

Darunavir binds predominantly to α1-acid glycoprotein. This protein binding is concentration-dependent indicative for saturation of binding. Therefore, there is a potential for protein displacement of medicines that are highly bound to α1-acid glycoprotein.

Life-threatening and fatal drug interactions have been reported in patients treated with colchicine and strong inhibitors of CYP3A and P-glycoprotein (P-gp). Darunavir with low-dose ritonavir must not be combined with medicines that are highly dependent on CYP3A for clearance and for which increased systemic exposure is associated with serious or life-threatening events (narrow therapeutic index) (see section 4.3).

A clinical study using a cocktail of medicines that are metabolised by cytochromes CYP2D9, CYP2C19 and CYP2D6 demonstrated an increase in exposure of CYP2C9 and CYP2C19 activity and inhibition of CYP2D6 activity in the presence of darunavir/ritonavir, which may be attributed to the presence of low-dose ritonavir. Co- administration of darunavir/ritonavir with medicines which are primarily metabolised by CYP2D6 (such as flecainide, propafenone and metoprolol) may result in increased concentrations of these medicines, which could increase or prolong their therapeutic effect and adverse reactions. Co-administration of darunavir/ritonavir and medicines primarily metabolised by CYP2C9 (such as warfarin) and CYP2C19 (such as methadone) may decrease systemic exposure to such medicines, which could decrease or shorten their therapeutic effect.

Although the effect on CYP2C8 has only been studied in vitro, co-administration of darunavir/ritonavir and medicines primarily metabolised by CYP2C8 (such as paclitaxel, rosiglitazone and repaglinide) may decrease systemic exposure to such medicines, which could decrease their therapeutic effect.

Ritonavir inhibits the transporter P-glycoproteins, OATP1B1 and OATP1B3, and co-administration with substrates of these transporters can increase plasma concentrations of these compounds (e.g. digoxin, statins, bosentan, and bosentan; see the table below).

Medicines that affect darunavir/ritonavir exposure
Darunavir and ritonavir are metabolised by CYP3A. Medicines that induce CYP3A activity are expected to increase the clearance of darunavir and ritonavir, resulting in lower plasma concentrations of these compounds, leading to loss of therapeutic effect and possible development of resistance (see sections 4.3). CYP3A inducers that are contraindicated include rifampicin, St John's wort and rifapentine.

Co-administration of darunavir/ritonavir with other medicines that inhibit CYP3A may decrease the clearance of darunavir and ritonavir, which may increase plasma concentrations of darunavir and ritonavir. Strong CYP3A inhibitors should be co-administered only if clinically vital and precautions taken to ensure effective levels of darunavir; these interactions are described in the table below (e.g. indinavir, systemic azoles like ketoconazole and clotrimazole).

Interaction table
Interactions between darunavir/ritonavir and antiretroviral and non-antiretroviral medicinal products are listed in the table below. The direction of the arrow for each pharmacokinetic parameter is based on the 90% confidence interval of the geometric mean ratio and is given by within (↔), below (↓) or above (↑) the 80-125% range.

Several interaction studies (indicated by # in the table below) used doses of darunavir that are lower than recommended or a different dosing regimen (see section 4.2 Posology). The effects on co-administered medicines may thus be underestimated, and clinical monitoring of safety may be indicated.

The below list of examples of drug-drug interactions is not comprehensive and therefore the product information for each drug that is co-administered with Darunavir and Ritonavir Tablets 400mg/50mg should be consulted for information on the route of metabolization, interaction pathways, potential risks, and specific actions to take with regard to co-administration.

Table : 1 Darunavir and Ritonavir Tablets 400mg/50mg Interactions

Medicines by therapeutic area	Interaction	Recommendations on co-administration
HIV ANTIRETROVIRALS		
Integrase strand transfer inhibitors		
Dolutegravir	Dolutegravir ↔ Dolutegravir AUC ↓ Cmax ↓	Darunavir/ritonavir co-administered with dolutegravir can be used without dose adjustment.
Raltegravir	Some clinical studies suggest raltegravir may decrease darunavir and ritonavir concentrations.	The effect of raltegravir on darunavir plasma concentrations is unknown. Darunavir/ritonavir does not appear clinically relevant. Darunavir/ritonavir and raltegravir can be used without dose adjustments.
Nucleos(t)ide reverse transcriptase inhibitors (NRTIs)		
Didanosine	Darunavir ↔ AUC ↔ Cmin ↔ Cmax ↔ didanosine AUC ↓ Cmax ↓	Darunavir/ritonavir and didanosine can be used without dose adjustments. Didanosine is to be taken on an empty stomach, thus it should be taken 1 hour before or 2 hours after darunavir/ritonavir given with food.
Tenofovir disoproxil	Darunavir AUC ↑ Cmin ↑ Cmax ↑ tenofovir AUC ↑ Cmax ↑ Cmin ↑ (Therefore from effect on MDR-1 transport in the renal tubules)	Monitoring of renal function may be indicated when darunavir/ritonavir is given in combination with tenofovir, particularly in patients with underlying systemic or renal disease, or in patients taking nephrotoxic agents.
Emtricitabine/ tenofovir alafenamide	Tenofovir alafenamide ↔ (≥ 200 mg/100 mg once daily)	The recommended dose of emtricitabine/tenofovir alafenamide when used with darunavir/ritonavir.
Abacavir/ Emtricitabine/ Lamivudine/ Stavudine/ Zidovudine	Not studied. Based on the elimination pathways of zidovudine, emtricitabine, stavudine and lamivudine, that are primarily renally excreted, and of abacavir for which metabolism is not mediated by CYP450, no interactions are expected with abacavir, and emtricitabine, lamivudine and stavudine/ritonavir.	Darunavir/ritonavir can be used with these NRTIs without dose adjustment.
Non-nucleos(t)ide reverse transcriptase inhibitors (NNRTIs)		
Etravirine	#Darunavir AUC ↓ Cmin ↓ Cmax ↓ etravirenz AUC ↑ Cmin ↑ Cmax ↑ (↑ effect/ risk from CYP3A inhibition of darunavir from CYP3A induction)	Clinical monitoring for central nervous system toxicity associated with increased exposure to etravirenz may be indicated when darunavir/ritonavir is given in combination with etravirenz. Etravirenz in combination with darunavir/ritonavir 800 mg/100 mg once daily may result in sub-optimal darunavir Cmin. If etravirenz is to be used in combination with darunavir, it should be dosed twice daily when boosted (see section 4.4). Darunavir/ritonavir and etravirine 200 mg twice daily can be used without dose adjustments.
Efavirenz	#Darunavir AUC ↓ Cmin ↓ Cmax ↓ efavirenz AUC ↑ Cmin ↑ Cmax ↑	Darunavir/ritonavir and efavirenz can be used without dose adjustments.
Nevirapine	#Darunavir: concentrations were expected to decrease nevirapine AUC ↑ Cmin ↑ Cmax ↑ (↑ nevirapine from CYP3A inhibition)	Darunavir/ritonavir and nevirapine can be used without dose adjustments.
Rilpivirine	Darunavir AUC ↔ Cmin ↓ Cmax ↓ rilpivirine AUC ↑ Cmin ↑ Cmax ↑	Darunavir/ritonavir and rilpivirine can be used without dose adjustments.
HIV Protease inhibitors (PIs) - without additional low-dose ritonavir		
Atazanavir	#Darunavir and ritonavir ↔ Cmin ↔ Cmax ↔ atazanavir AUC ↔ Cmin ↑ Cmax ↓	Darunavir/ritonavir and atazanavir can be used without dose adjustments.
Indinavir	#Darunavir AUC ↑ Cmin ↑ Cmax ↑ indinavir AUC ↑ Cmin ↑ Cmax ↔	When used in combination with darunavir/ritonavir, the dose of indinavir may be reduced from 800 mg twice daily to 600 mg twice daily to manage side effects.
Saquinavir	#Darunavir AUC ↓ Cmin ↓ Cmax ↓ saquinavir AUC ↑ Cmin ↑ Cmax ↓	It is not recommended to combine darunavir/ritonavir with saquinavir.
HIV Protease inhibitor (PI) – with additional low dose ritonavir		
Lopinavir/ritonavir 400 mg/100 mg twice daily	Darunavir AUC ↓ Cmin ↓ Cmax ↓ lopinavir AUC ↑ Cmin ↑ Cmax ↓	Due to a decrease in the exposure of darunavir by 40%, appropriate doses of the combination have not been established. Hence, concomitant use of darunavir/ritonavir and the combination product lopinavir/ritonavir is contraindicated (see section 4.3).
CCR5 ANTAGONIST		
Maraviroc	Darunavir, ritonavir concentrations were consistent with historical data maraviroc AUC ↑ Cmax ↑	Maraviroc dose should be 150 mg twice daily when co-administered with darunavir/ritonavir
α1-ADRENORECEPTOR ANTAGONIST		
Alfuzosin	Darunavir/ritonavir is expected to increase alfuzosin plasma concentrations (CYP3A inhibition).	Co-administration of darunavir/ritonavir and alfuzosin is contraindicated (see section 4.3).

ANTIANGINA/ANTIARRHYTHMICS		
Disopyramine Flecainide Lidocaine(systemic)Mexiletine Propafenone Amiodarone Bepridil Dronedarone hydrazine Quinidine Ranolazine	Not studied. Darunavir/ritonavir is expected to increase plasma concentrations of these medicines. (CYP3A and CYP2D6 inhibition)	Caution is warranted and therapeutic concentration monitoring, if available, is recommended for these medicines when co-administered with darunavir/ritonavir. Darunavir/ritonavir co-administration with amiodarone, bepridil, dronedarone, ivabradine, quinidine, or ranolazine is contraindicated (see section 4.3).
Digoxin	Digoxin AUC ↑ Cmax ↑ (↑ digoxin from probable inhibition of P-gp)	Because digoxin has a narrow therapeutic index, it is recommended that the lowest possible dose of digoxin should initially be prescribed for patients taking darunavir/ritonavir. The digoxin dose should be carefully titrated to obtain the desired clinical effect while assessing the overall clinical state of the subject.
ANTIBIOTICS		
Clarithromycin	#Darunavir AUC ↓ Cmin ↑ Cmax ↓ clarithromycin AUC ↑ Cmin ↑ Cmax ↑ (↑ clarithromycin from CYP3A inhibition and possible P-gp inhibition)	Caution should be exercised when clarithromycin is combined with darunavir/ritonavir. For patients with renal impairment the product information of clarithromycin should be consulted for the recommended dose.
Rifamycins		
Rifampicin Rifapentine	Not studied. Rifapentine and rifampicin are strong CYP3A inducers and decrease concentrations of other protease inhibitors, which can result in virological failure and development of resistance. Moreover, concomitant administration of rifampicin has been reported to cause severe hepatocellular reactions when used with rifampicin.	The combination of rifampicin and darunavir/ritonavir is contraindicated (see section 4.3). The combination of rifapentine and darunavir/ritonavir is not recommended.
Rifabutin	Darunavir AUC ↑ Cmin ↑ Cmax ↑ rifabutin AUC ↑ Cmax ↔ (Rifabutin is an inducer and substrate of CYP3A.)	A dosage reduction of rifabutin by 75% of the usual dose of 300 mg/day (to 150 mg once every other day) and increased monitoring for rifabutin-related adverse events is warranted in patients receiving darunavir/ritonavir. In case of side effects, a further increase of the dosing interval for rifabutin and monitoring of rifabutin levels should be considered. Based on the safety profile of darunavir/ritonavir, the increase in darunavir exposure in the presence of rifabutin does not warrant a dose adjustment for darunavir/ritonavir.
ANTICOAGULANTS / PLATELET AGGREGATION INHIBITORS		
Apixaban Edoxaban Rivaroxaban	Not studied. Co-administration of darunavir/ritonavir with these anticoagulants may increase concentrations of the anticoagulant and increase bleeding risk. (CYP3A and P-gp inhibition)	The use of darunavir/ritonavir and these anticoagulants is not recommended.
Dabigatran Ticagrelor	Not studied. Co-administration with darunavir/ritonavir is expected to substantially increase exposure to dabigatran or ticagrelor.	Concomitant administration of darunavir/ritonavir with dabigatran or ticagrelor is contraindicated (see section 4.3).
Clopidogrel	Not studied. Co-administration of clopidogrel with darunavir/ritonavir is expected to decrease plasma concentrations of clopidogrel active metabolite, which may reduce the antiplatelet activity of clopidogrel.	Co-administration of clopidogrel with darunavir/ritonavir is not recommended. Use of antiplatelets not affected by CYP inhibition or induction (e.g. prasugrel) is recommended.
ANTICONVULSANTS		
Phenobarbital Phenytoin	Not studied. Phenobarbital and phenytoin are expected to decrease plasma concentrations of darunavir/ritonavir (induction of CYP450 enzymes).	Darunavir/ritonavir should not be used in combination with these medicines.
Carbamazepine	Not studied. Carbamazepine AUC ↓ Cmin ↓Cmax ↓ carbamazepine AUC ↑ Cmin ↑ Cmax ↑	No dose adjustment for darunavir/ritonavir is recommended. If darunavir/ritonavir and carbamazepine need to be co-administered, patients should be monitored for carbamazepine related adverse events. Carbamazepine concentrations should be monitored if possible and its dose titrated for adequate response. The carbamazepine dose should be reduced by as much as 25-50% in the presence of dose-stabilized darunavir/ritonavir.
Clonazepam	Not studied. Co-administration of darunavir/ritonavir with clonazepam may increase concentrations of clonazepam. (CYP3A inhibition)	Clinical monitoring is recommended when co-administering darunavir/ritonavir and clonazepam.
ANTIDEPRESSANTS		
Paroxetine	#Darunavir AUC ↔ Cmin ↔ Cmax ↔ paroxetine AUC ↓ Cmin ↓ Cmax ↓	If antidepressants are co-administered with darunavir/ritonavir, dose titration of the antidepressant based on an assessment of antidepressant response is recommended. Also, patients on a stable dose of antidepressants who start treatment with darunavir/ritonavir should be monitored for antidepressant response.
Sertraline	#Darunavir AUC ↔ Cmin ↔ Cmax ↔ sertraline AUC ↔ Cmin ↔ Cmax ↔	Darunavir/ritonavir can be used without dose adjustments.
Amitriptyline Desipramine Imipramine Nortriptyline Trazodone	Concomitant use of darunavir/ritonavir and these antidepressants may increase concentrations of the antidepressant. (CYP2D6 and CYP3A inhibition)	Clinical monitoring is recommended when co-administering darunavir/ritonavir with these antidepressants and dose adjustment of the antidepressant may be needed.
ANTIEMETICS		
Domperidone	Not studied. Domperidone is mainly metabolised by CYP3A.	Co-administration of domperidone with darunavir/ritonavir is contraindicated because risk of domperidone's cardiac adverse events may be increased.
ANTIFUNGALS		
Voriconazole	Not studied. Ritonavir may decrease plasma concentrations of voriconazole. (induction of CYP450 enzymes)	Voriconazole should not be combined with darunavir/ritonavir unless an assessment of the benefits and risks justifies the use of voriconazole.
Ketoconazole	#Darunavir AUC ↑ Cmin ↑ Cmax ↑ ketoconazole AUC ↑ Cmin ↑ Cmax ↑ (CYP3A inhibition)	Caution is warranted and clinical monitoring is recommended. When co- administration is required the daily dose of ketoconazole should not exceed 200 mg.
Fluconazole Isavuconazole Itraconazole Posaconazole	Not studied. Darunavir may increase antifungal plasma concentrations. Isavuconazole, Itraconazole, or posaconazole may increase darunavir concentrations. (CYP3A inhibition and P-gp inhibition)	Caution is warranted and clinical monitoring is recommended. When co- administration is required the daily dose of fluconazole should not exceed 200 mg.
Clotrimazole	Not studied. Concomitant systemic use of clotrimazole and darunavir/ritonavir is expected to decrease plasma concentrations of darunavir/ritonavir due to CYP3A inhibition. darunavir AUC/24h ↑ (based on population pharmacokinetic model)	Caution is warranted and clinical monitoring is recommended. when co- administration of clotrimazole is required.
ANTIGOUT MEDICINES		
Colchicine	Not studied. Concomitant use of colchicine and darunavir/ritonavir may increase the exposure to colchicine. (CYP3A and P-gp inhibition).	Reducing colchicine dosage or interrupting colchicine treatment is recommended in patients with normal renal and hepatic function if treatment with darunavir/ritonavir is required. Patients with renal or hepatic impairment must not be given colchicine with darunavir/ritonavir (see sections 4.3 and 4.4).
ANTIMALARIALS		
Artemether/lumefantrine	Darunavir AUC ↔ Cmin ↓ Cmax ↔ artemether and dihydroartemisinin AUC ↓ Cmin ↔↓ Cmax ↓ lumefantrine AUC ↑ Cmin ↑ Cmax ↑	The combination of darunavir and artemether/lumefantrine is contraindicated, due to the increase in lumefantrine exposure.
Artemisinin	Coadministration may increase comedication exposure, and a dose adjustment may be needed. Monitor clinical effect.	Dose adjustment may be needed. Monitor clinical effect.
Halofantrine	Not studied. Halofantrine is extensively metabolized by CYP3A4 Inhibition of halofantrine metabolism by ritonavir is expected to increase halofantrine concentrations, potentially prolong the QT interval.	Concomitant administration of boosted darunavir and halofantrine is not recommended. Halofantrine has a narrow therapeutic index with an increased risk of QT- prolongation at higher exposures.
Mefloquine	Coadministration may increase comedication exposure.	Caution and dose monitoring is recommended.
Proguanil	Coadministration may decrease proguanil exposure.	Coadministration of atovaquone/proguanil should be avoided whenever possible. If judged clinically necessary, consider taking atovaquone/proguanil with a high fat meal to increase its bioavailability and increase the dosage if required.
Quinine	Coadministration may increase quinine exposure.	In addition, patients should be counselled as quinine has a risk of QT prolongation. EGG monitoring is recommended.
ANTINEOPLASTICS		
Dasatinib Nilotinib Vinblastine Vincristine	Not studied. Darunavir/ritonavir is expected to increase plasma concentrations of these antineoplastic medicines. (CYP3A inhibition)	Concentrations of these medicines may increase when co-administered with darunavir/ritonavir, potentially resulting in increased adverse events usually associated with these agents. Caution should be exercised when combining these antineoplastic medicines with darunavir/ritonavir Concomitant use of vinblastine or vincristine and darunavir/ritonavir is not recommended.
CORTICOSTEROIDS		
Corticosteroids primarily metabolised by CYP3A (including betamethasone, budesonide, fluticasone, mometasone, prednisone, dexamethasone (systemic))	Darunavir AUC ↓ Cmin ↓ Cmax ↓ fluticasone propionate AUC ↑ Cmin ↑ Cmax ↑ Other corticosteroids: Concentrations of these medicines may increase when co-administered with darunavir/ritonavir, resulting in reduced serum cortisol concentrations	Concomitant administration of darunavir/ritonavir and corticosteroids dependent by CYP3A (e.g. fluticasone propionate or other inhaled or nasal corticosteroids) may increase the risk of systemic corticosteroid effects, including Cushing's syndrome and adrenal suppression. Co-administration with these corticosteroids is not recommended unless the potential benefit of treatment outweighs the risk, in which case patients should be monitored for systemic corticosteroid effects. Moreover, in case of withdrawal of corticosteroids, progressive dose reduction may have to be considered.
ENDOTHELIN RECEPTOR		
Bosentan	Not studied. Concomitant use of bosentan and darunavir/ritonavir may increase plasma concentrations of bosentan. Bosentan is expected to decrease plasma concentrations of darunavir/ritonavir (CYP3A induction).	When administered concomitantly with darunavir/ritonavir, the patient should be monitored for bosentan side effects.
HEPATITIS C VIRUS DIRECT-ACTING ANTIVIRALS		
Elbasvir/grazoprevir	Darunavir/ritonavir may increase the exposure to grazoprevir and elbasvir (CYP3A and OATP1B inhibition)	Concomitant use of darunavir/ritonavir and elbasvir/grazoprevir is contraindicated (see section 4.3).
Glecaprevir/pibrentasvir	Darunavir/ritonavir may increase the exposure to glecaprevir and pibrentasvir (P-gp, BCRP and OATP1B1/3 inhibition)	Co-administration of darunavir/ritonavir with glecaprevir/ pibrentasvir is not recommended.
Ombitasvir/ paritaprevir/ ritonavir	Darunavir and Ritonavir AUC ↓ Cmin ↓ Cmax ↑ ombitasvir Cmax ↔ AUC ↔ Cmin ↔ paritaprevir Cmax AUC Cmin ↑	Darunavir and Ritonavir Tablets 400mg/50mg should not be used because the co-administered medicine already contains ritonavir.
Ombitasvir/paritaprevir/ ritonavir	Darunavir Cmax ↔ AUC ↔ Cmin ↓ ombitasvir Cmax ↔ AUC ↔ Cmin ↔ paritaprevir Cmax ↑ AUC ↑ Cmin ↑	
Ledipasvir	Darunavir Cmax ↔ AUC ↔ Cmin ↔ ledipasvir Cmax ↑ AUC ↑ Cmin ↑	No dose adjustment is required.
Sofosbuvir	darunavir Cmax ↔ AU C ↔ Cmin ↔ sofosbuvir Cmax ↑ AUC ↑ GS-331007 Cmax ↔ AUC ↔	
Daclatasvir	Darunavir AUC ↔ Cmax ↔ Cmin ↔ daclatasvir AUC ↔ Cmax ↔	No dose adjustment is required.
HERBAL PRODUCTS		
St. John's wort (*Hypericum perforatum*)	Not studied. St. John's wort is expected to decrease the plasma concentrations of darunavir and ritonavir (CYP450 induction).	Darunavir/ritonavir must not be used concomitantly with products containing St John's wort (see section 4.3). If a patient is already taking St. John's wort, stop St John's wort and if possible check viral levels. Darunavir (and ritonavir) exposure may increase on stopping St John's wort. The induction effect may persist for at least 2 weeks after stopping St John's wort.
HMG CO-A REDUCTASE INHIBITORS		
Lovastatin Simvastatin	Not studied. Lovastatin and simvastatin AUC ↑↑ are substantially increased when co-administered with darunavir/ritonavir. (CYP3A inhibition)	Increased plasma concentrations of lovastatin or simvastatin may cause myopathy, including rhabdomyolysis. Concomitant use of darunavir/ritonavir, with lovastatin and simvastatin is therefore contraindicated (see section 4.3).
Atorvastatin	Atorvastatin AUC ↑ Cmin ↑ Cmax ↑	When administration of atorvastatin and darunavir/ritonavir is desired, it is recommended to start with an atorvastatin dose of 10 mg once daily. A gradual dose increase of atorvastatin may be tailored to the clinical response.
Pravastatin	Pravastatin AUC ↑ Cmax ↑	When administration of pravastatin and darunavir/ritonavir is required, it is recommended to start with the lowest possible dose of pravastatin and titrate up to the desired clinical effect while monitoring for safety.
Rosuvastatin	Rosuvastatin AUC ↑

1. NOME DO MEDICAMENTO | **POM** | **Schedule: S2** | **NS2** | **PP**

Darunavir e Ritonavir Comprimidos 400 mg/80 mg | **Medicamento apenas com receita - Lista I**

2. COMPOSIÇÃO QUALITATIVA E QUANTITATIVA

Cada comprimido revestido por película contém 433,640 mg de Etanolato de Darunavir equivalente a 400 mg de Darunavir e Ritonavir USP 80 mg.

Lista completa de excipientes, vide seção 6.1

3. FORMA FARMACÊUTICA

Comprimidos Revestidos por Película

Comprimidos revestidos por película amarelos, em forma de cápsula, biconvexos, com bordas chanfradas, gravados com 'H' em um lado e '136' no outro lado.

4. Informações Clínicas

4.1 Indicações terapêuticas

Darunavir e Ritonavir Comprimidos 400 mg/80 mg é indicado em combinação com outros medicamentos antiretrovirais para o tratamento da infecção pelo vírus da imunodeficiência humana tipo 1 (HIV-1) em adultos e adolescentes com peso mínimo de 40 kg que:

- não foram previamente tratados com terapia antirretroviral (pacientes virgens de tratamento à protease inibidor), vide seção 4.2); ou
- foram previamente tratados com um inibidor da protease ('experiência com inibidor da protease'), mas não apresentam mutações associadas à resistência ao darunavir (vide abaixo) e que têm RNA do HIV-1 no plasma < 100 000 cópias/ml e contagem de células CD4+ > 100 × 106 células/l, (vide seções 4.4 e 5.1).

Darunavir e Ritonavir Comprimidos 400 mg/80 mg é também para ser experiência com inibidores de protease e tenho de genótipo HIV-1 não estiver disponível. As mutações associadas à resistência ao Darunavir incluem: V11I, V32I, L33F, I47V, I50V, I54M, I54L, T74P, L76V, I84V e L89V.

Ver a seção 4.4 para aconselhamento sobre o tratamento com Darunavir/ritonavir em doentes com experiência com inibidores da protease quando o teste de genótipo não está disponível.

As diretrizes oficiais para o tratamento de pessoas infectadas pelo HIV-1 (p.ex., as do DAWG) devem ser consideradas.

4.2 Posologia e Modo de Administração

A terapia deve ser iniciada por um profissional de saúde com experiência no tratamento da infecção pelo HIV.

Posologia

Pacientes com peso de pelo menos 40 kg

Em pacientes virgens de IP e em pacientes com experiência de IP sem mutações associadas à resistência ao darunavir e que têm RNA do HIV-1 plasmático < 100 000 cópias/ml e contagem de células CD4+ ≥ 100 × 106 células/l, (vide seção 4.1), a dose recomendada de Darunavir e Ritonavir Comprimidos 400 mg/80 mg é:

2 comprimidos tomados uma vez ao dia com refeição, por volta da mesma hora todos os dias.

Darunavir e Ritonavir Comprimidos 400 mg/80 mg não são apropriados para o tratamento de todos os outros pacientes com experiência em TARV ou se teste do genótipo HIV-1 não estiver disponível.

Dose pediátrica

Se uma dose de Darunavir e Ritonavir Comprimidos 400 mg/80 mg for omitida dentro de 12 horas do horário habitualmente tomado, os pacientes devem ser instruídos a tomar a dose de Darunavir e Ritonavir Comprimidos 400 mg/80 mg com refeição o mais rápido possível. Se tiverem passado mais de 12 horas após a hora habitual de administração, a dose em falta não deve ser tomada e o doente deve retomar o esquema posológico habitual.

Se um paciente vomitar dentro de 4 horas após tomar Darunavir e Ritonavir Comprimidos 400 mg/80 mg, o paciente deve tomar outra dose com uma refeição o mais rápido possível. Se o paciente vomitar mais de 4 horas após tomar o medicamento, o paciente não precisa de tomar outra dose.

Populações especiais

Idosos

A informação disponível nessa população é limitada e, por conseguinte, Darunavir e Ritonavir comprimidos 400 mg/80 mg devem ser utilizados com precaução nesse grupo etário (vide seções 4.4 e 5.2).

Disfunção hepática

Darunavir é metabolizado pelo sistema hepático. Não se recomenda a quota de dose em pacientes com disfunção hepática leve (Child-Pugh Classe A) ou moderada (Child-Pugh Classe B); entretanto, Darunavir e Ritonavir Comprimidos 400 mg/80 mg devem ser usados com cautela nesses pacientes. Não existem dados em pacientes com disfunção hepática grave. A disfunção hepática grave pode resultar em aumento da exposição ao darunavir e pois dos efeitos colaterais. Assim, Darunavir e Ritonavir comprimidos 400 mg/80 mg não devem ser utilizados em doentes com disfunção hepática grave (Child-Pugh Classe C) (vide seções 4.3, 4.4 e 5.2).

Disfunção renal

Não é necessário ajuste da dose para Darunavir e Ritonavir Comprimidos 400 mg/80 mg em pacientes com disfunção renal (vide seções 4.4 e 5.2).

População pediátrica

Darunavir e Ritonavir Comprimidos 400 mg/80 mg não é adequado para crianças com peso inferior a 40 kg. Poderão ser necessárias outras formulações de Darunavir/ritonavir.

Darunavir/ritonavir não deve ser utilizado em crianças com menos de 3 anos de idade ou com peso inferior a 15 kg (vide seção 5.3).

Gravidez e pós-parto

Durante a gravidez e o pós-parto não é necessário ajuste de dose para Darunavir e Ritonavir (vide seções 4.6 e 5.2).

Modo de administração

Os pacientes devem ser instruídos a tomar darunavir/ritonavir dentro de 30 minutos após uma refeição. O tipo de alimento não afeta a exposição ao darunavir (vide seção 5.2).

4.3 Contra-indicações

Hipersensibilidade à substância ativa ou a qualquer um dos excipientes listados na seção 6.1 Pacientes com disfunção hepática grave (Classe C de Child-Pugh).

Tratamento concomitante com os medicamentos listados abaixo porque podem diminuir as concentrações plasmáticas de darunavir e ritonavir, o que pode levar à perda do efeito terapêutico e desenvolvimento de resistência (vide seções 4.4 e 4.5). Os seguintes não devem ser usados com Darunavir e Ritonavir Comprimidos 400 mg/80 mg:

- lopinavir/ritonavir,
- os indutores fortes do CYP3A, rifampicina e preparados à base de planta contendo erva de São João (*Hypericum perforatum*).

Darunavir potencializ com ritonavir inibe a eliminação de substâncias ativas que são altamente dependentes do CYP3A para depuração. Estas substâncias ativas concomitantes com medicamentos para os quais as concentrações plasmáticas elevadas estão associadas a efeitos colaterais graves ou com ocorrência de reação contra-indicada (vide também seção 4.5). Essas substâncias ativas incluem:

- alfuzosina,
- amiodarona, bepridil, dronedarona, encainida, flecainida, ivabradina, propafenona, quinidina, ranolazina
- astemizol, terfenadina
- avanafil, sildenafil quando usado para o tratamento da hipertensão arterial pulmonar, vardenafil
- cisaprida
- clorzoxazol, diazepam, estazolam, flurazepam, midazolam oral e triazolam
- ciclospo, lurasidona, pimozida, quetiapina, sertindol
- colchicina quando usado em pacientes com disfunção renal ou hepática
- dabigatrana, fusidato
- dipiridamol
- disopiramida
- derivados do ergot (p.e., diidroergocristina, ergometrina, ergotamina, metilergonovina)
- eliglustr/ginsgrenir, ombitasvir/paritaprevir/ritonavir
- ácido fusídico
- halofantrina, lumefantrina
- lomitapida, tovaptidina e simvastatina
- nalmefen
- nevirapina, vandetanib
- petidina, piroxicam, propoxifeno

4.4 Advertências e precauções especiais de utilização

Pacientes com experiência em inibidores de protease

Quando o teste genotípico não é viável em pacientes com experiência com inibidores de protease, recomenda-se darunavir/ritonavir 600 mg/100 mg duas vezes ao dia. Outras formulações devem ser usadas; Darunavir e Ritonavir Comprimidos 400 mg/80 mg não é adequado.

Transmissão

Embora a supressão viral eficaz com terapia antirretroviral reduz substancialmente o risco de transmissão sexual, pode permanecer um risco residual. As precauções para prevenir a transmissão devem ser tomadas de acordo com as diretrizes nacionais.

Aconselha-se a avaliação regular da resposta virológica. No cenário de falha ou perda de resposta virológica, deve ser realizado o teste de resistência.

Pacientes com experiência em terapia antirretroviral – dose única diária

Não recomendamos Darunavir e Ritonavir 800 mg/100 mg uma vez ao dia em pacientes com experiência em TARV não deve ser usado em pacientes com uma ou mais mutações associadas à resistência ao darunavir (DRV-RAMs) ou RNA do HIV-1 ≥ 100 000 cópias/ml ou contagem de células CD4+ < 100 × 106 células/l. Dados limitados em pacientes com experiência com esquema de dose única não nucleotídeos (ou nucleosídeos) inibidores da transcriptase reversa (NRTIs) foram estudados nessa população. Os dados são limitados em doentes com subtipos de HIV-1 diferentes de B (vide seção 5.1).

Idosos

Como as informações sobre o uso de darunavir/ritonavir em pacientes com 65 anos ou mais são limitadas, Darunavir e Ritonavir comprimidos 400 mg/80 mg devem ser usados com cautela em pacientes desse grupo etário, refletindo a maior frequência de diminuição da função hepática e de doença concomitante ou outra terapia (vide seções 4.2 e 5.2).

Reações cutâneas graves

Foram relatadas reações cutâneas graves, que podem ser acompanhadas de febre e elevação das transaminases. DRESS (eruption cutânea com eosinofilia e sintomas sistémicos) e síndrome de Stevens-Johnson relatados ocasionalmente (< 0,1%) e durante a experiência pós-comercialização foram relatadas necrólise epidérmica tóxica e pustulose exantemática generalizada aguda. Os sintomas podem incluir erupção cutânea ou reações associadas a vários órgãos (p.e., hepatite, eosinofilia, exantema, angioedema, mal-estar, síndrome de Stevens-Johnson, cansaço, conjuntivite, hepatite e eosinofilia (vide também seção 4.8). Darunavir e Ritonavir comprimidos 400 mg/50 mg devem ser descontinuados imediatamente se surgirem sinais ou sintomas de reações cutâneas graves.

A erupção cutânea ocorreu mais frequentemente em doentes com experiência de tratamento a receber regimes contendo darunavir/ritonavir + raltegravir comparativamente a doentes a receber darunavir/ritonavir sem raltegravir ou raltegravir sem darunavir (vide seção 4.8).

Darunavir contém uma porção de sulfonamida. Darunavir e Ritonavir comprimidos 400 mg/50 mg devem ser usados com cautela em pacientes com alergia a sulfonamidas.

Hepatotoxicidade

Hepatite induzida por substâncias (p.e., hepatite aguda, hepatite citolítica) foi relatada em 0,5% dos pacientes recebendo terapia antirretroviral combinada com Darunavir/ritonavir. Os doentes com disfunção hepática, incluindo hepatite B ou C crónica ativa, têm um risco aumentado de anomalias da função hepática, incluindo reações adversas hepáticas graves e potencialmente fatais. Em caso de terapia antiviral concomitante para hepatite B ou C, consulte também as informações do produto relevante para esses medicamentos.

Devem ser realizadas os testes laboratoriais apropriados antes de iniciar a terapia com darunavir/ritonavir e os pacientes devem ser monitorizados durante o tratamento. O aumento no comprimento de ALT/AST deve ser considerado em pacientes com hepatite crónica subjacente, cirrose ou em pacientes que experience elevações de transaminases pré-tratamento, especialmente durante os primeiros meses de tratamento com darunavir/ritonavir. Foi observar evidência de disfunção hepática nova para (incluindo elevação clinicamente significativa das enzimas hepáticas e/ou sintomas como fadiga, anorexia, náuseas, icterícia, urina escura, sensibilidade hepática, hepatomegalia) em pacientes usando darunavir/ritonavir, é interrupção do tratamento deve ser considerada prontamente.

Disfunção hepática

A segurança e eficácia de darunavir não foram estabelecidas em pacientes com distúrbios hepáticos graves o Darunavir e Ritonavir comprimidos 400 mg/80 mg é, portanto, contra-indicado em pacientes com disfunção hepática grave. Devido a um aumento nas concentrações plasmáticas de darunavir em doentes com disfunção hepática moderada, este medicamento deve ser utilizado com precaução em doentes com disfunção hepática ligeira ou moderada (vide seções 4.2, 4.3 e 5.2).

Disfunção renal

Não são necessárias precauções especiais ou ajuste de dose para Darunavir e Ritonavir Comprimidos 400 mg/50 mg em pacientes com disfunção renal. Como o darunavir e o ritonavir se ligam fortemente às proteínas plasmáticas, é improvável que sejam significativamente removidos por hemodiálise ou diálise peritoneal. Portanto, não são necessárias precauções especiais ou ajustes de dose nesses doentes (vide seções 4.2 e 5.2).

Suspeita uma hemofilia

Houve relatos de aumento de sangramento, incluindo hematomas cutâneos espontâneos e hemartrose em pacientes com hemofilia tipo A e B tratados com inibidores de protease. Em alguns pacientes foi administrado fator VIII adicional. Em mais da metade dos casos relatados, o tratamento com inibidores de protease foi continuado ou reintroduzido se o tratamento tivesse sido descontinuado. Foi sugerida uma relação causal, mas o mecanismo de ação não foi elucidado. Os pacientes com hemofilia devem, portanto, ser alertados sobre a possibilidade de aumento do sangramento.

Peso, lipídios e glicose no sangue

O peso e nos níveis de lípidos e glicose no sangue podem aumentar durante a terapia antirretroviral. Tais mudanças podem, em parte, estar ligadas ao controle da doença e ao estilo de vida. Para os lípidos, existem evidências de um efeito do tratamento, enquanto para o ganho de peso não existem evidências fortes relacionando com qualquer tratamento específico. Para a monitorização dos lípidos e glicose sanguíneos consulte as diretrizes de tratamento do HIV estabelecidas. Os distúrbios lipídicos devem ser tratados conforme clinicamente apropriado.

Osteonecrose

Foram relatados casos de osteonecrose particularmente em pacientes com doença avançada por HIV ou exposição prolongada à terapia antirretroviral combinada. A etiologia é considerada multifatorial (incluindo uso de corticosteroides, consumo de álcool, imunossupressão grave, índice de massa corporal elevado). Os doentes devem ser aconselhados a procurar aconselhamento médico se tiverem artralgias e dores nas articulações, rigidez nas articulações ou dificuldade de movimento.

Infecções oportunistas

Os pacientes que recebem darunavir ou qualquer outra terapia antirretroviral podem continuar a desenvolver infecções oportunistas e outras complicações da infeção do HIV. Portanto, os pacientes devem permanecer sob observação clínica frequente por profissionais de cuidados de saúde experientes nessas doenças associadas à infeção pelo HIV.

Síndrome inflamatória de reconstituição imune

No início da terapia antirretroviral combinada (TARC) em pacientes com deficiência imune grave, pode ocorrer uma reação inflamatória a infeções oportunistas assintomáticas ou residuais e pode causar condições clínicas graves ou agravar os sintomas. Normalmente, essas reações ocorrem nas primeiras semanas ou meses após o início da TARC. Exemplos relevantes são retinite por citomegalovírus, infeções micobacterianas generalizadas e/ou focais e pneumonia causada por *Pneumocystis jirovecii* (anteriormente conhecida como *Pneumocystis carinii*). Qualquer sintoma inflamatório deve ser avaliado e tratado quando necessário. Além disso, ocorreu melhoria de terapias simples e herpes zoster em reações ativadas após o início do tratamento.

Também foram relatados doenças autoimunes (como a doença de Graves e hepatite autoimune) no cenário de reconstituição imune; porém, o tempo de início é mais variável e esses eventos podem ocorrer muitos meses após o início do tratamento. (vide seção 4.8).

Interações com medicamentos

Efavirenz em combinação com Darunavir e Ritonavir comprimidos 400 mg/50 mg uma vez ao dia pode resultar em concentração de darunavir abaixo do ideal. Se for necessário utilizar efavirenz, recomenda-se darunavir/ritonavir 600 mg/100 mg duas vezes por dia em doentes com experiência de inibidores da protease (vide seção 4.5).

Foram relatadas interações medicamentosas fatais e com risco de vida em pacientes tratados com colchicina e foram inibidores de CYP3A e glicoproteína-P (P-gp; vide seções 4.3 e 4.5).

O darunavir liga-se predominantemente à α-1-glicoproteína ácida. Essa ligação à proteína depende da concentração sendo indicativa de saturação da ligação. Portanto, existe um potencial para deslocamento de proteínas de medicamentos que se ligam fortemente à α1-glicoproteína ácida.

Foram relatadas interações medicamentosas fatais e com risco de vida em pacientes tratados com colchicina e fortes inibidores de CYP3A e glicoproteína-P (P-gp). O darunavir com ritonavir deve ser usado com cautela em combinação com medicamentos substratos dessas enzimas. Recomenda-se a redução da dose da colchicina ou a interrupção do tratamento com colchicina em pacientes em risco de disfunção renal ou hepática normais se o tratamento com darunavir/ritonavir for necessário. Os doentes com comprometimento renal ou hepático não devem receber colchicina com darunavir/ritonavir (vide também seções 4.3 e 4.4).

Um estudo clínico usando um cocktail de medicamentos que são metabolizadas pelo citocromo P450, CYP2C19 e CYP2D6 demonstrou que a administração de 600 mg/100 mg de darunavir duas vezes ao dia não teve efeito significativo sobre o CYP2C19, o que pode ser indicado a presença de efeito sobre o CYP2C19 em pacientes com medicamentos que são principalmente metabolizados pelo CYP2D6 (tais como flecainida, propafenona e metoprolol) pode resultar no aumento das concentrações plasmáticas destes medicamentos, a que pode aumentar ou prolongar o seu efeito terapêutico e as reações adversas. A coadministração de darunavir/ritonavir e medicamentos metabolizados principalmente pelo CYP2D6 (como a varfarina) e CYP2C19 (como a metadona) pode diminuir a exposição a esses medicamentos, o que pode diminuir os seus efeitos terapêuticos.

Embora o efeito sobre o CYP2D6 só tenha sido estudado in vitro, a coadministração de darunavir/ritonavir e medicamentos primariamente metabolizados pelo CYP2D6 como quetiapina, risperidona e a maioria dos inibidores seletivos da recaptação da serotonina (ISRSs), que pode diminuir ou encurtar o seu efeito terapêutico.

Ritonavir inibe as transportadoras de glicoproteína-P, OATP1B1 e OATP1B3, e a coadministração com substratos destas transportadoras pode aumentar as concentrações plasmáticas destes compostos (p.e., ebastato de dabigatrano, digoxina, estatinas e bosentano; vide tabela abaixo).

Medicamentos que afetam a exposição ao darunavir/ritonavir

O darunavir e o ritonavir são metabolizados pelo CYP3A. Prevê-se que os medicamentos que induzem a atividade do CYP3A aumentem a depuração de darunavir e ritonavir, resultando em concentrações plasmáticas mais baixas de darunavir e ritonavir. Prevê-se que os inibidores do CYP3A diminuam a depuração de darunavir e ritonavir, resultando em concentrações plasmáticas mais elevadas de darunavir e ritonavir. Os estudos de interação medicamentosa foram realizados com darunavir/ritonavir 600 mg/100 mg duas vezes por dia, a dose recomendada em combinação com outros medicamentos que são principalmente metabolizados pelo CYP3A (tais como fluconazol, itraconazol e metronidazol) pode resultar no aumento das concentrações plasmáticas de darunavir/ritonavir e medicamentos metabolizados principalmente pelo CYP3A (como a varfarina) e CYP2C19 (como a metadona) pode diminuir a exposição a esses medicamentos, o que pode diminuir os seus efeitos terapêuticos.

A coadministração de darunavir/ritonavir com outros medicamentos que induzem o CYP3A pode diminuir a depuração de darunavir e ritonavir, o que pode aumentar as concentrações plasmáticas de darunavir e ritonavir. Os indutores do CYP3A devem ser coadministrados somente se clinicamente vitais e devem ser tomadas precauções para garantir níveis efetivos de darunavir; essas interações são descritas na tabela abaixo (p.e., rifampicina, ácidos sistémicos corticosteroides e cimetidina).

Tabela de interações

As interações entre darunavir/ritonavir e medicamentos antiretrovirais e não antiretrovirais estão listadas na tabela abaixo. A direção da seta para cada parâmetro farmacocinético baseia-se no intervalo de confiança de 90% da razão da média geométrica dentro de (↑, abaixo (↓) ou dentro (↑↓) da faixa de 80-125%.

Vários estudos de interação (indicados por # na tabela abaixo) utilizaram doses de darunavir inferiores às recomendadas ou um regime posológico diferente (vide seção 4.2 Posologia). Os efeitos dos medicamentos coadministrados podem, portanto, ser subestimados e a monitorização clínica da segurança pode ser indicado.

A lista abaixo de exemplos de interações medicamentosas não é abrangente e, portanto, devem ser consultadas as informações do produto para cada medicamento coadministrado com Darunavir e Ritonavir Comprimidos 400 mg/50 mg para obter informações sobre a via de metabolismo, vias de interação, vias potenciais, e doenças específicas a serem tomadas em relação à coadministração.

Tabela : 1 Darunavir e Ritonavir Comprimidos 400 mg/50 mg

Medicamentos	Alterações	Recomendações sobre a coadministração
ANTIRETROVIRAIS DO HIV		
Inibidores da transferência da cadeia da integrase		
Dolutegravir	Darunavir ↔ Dolutegravir AUC ↓ Cmáx ↓	Darunavir/ritonavir coadministrado com dolutegravir pode ser usado sem ajuste de dose.
Raltegravir	Alguns estudos clínicos sugerem que o raltegravir pode diminuir plasmáticas de darunavir.	O efeito de raltegravir nas concentrações plasmáticas de darunavir não parece clinicamente relevante. Darunavir/ritonavir e raltegravir podem ser usados sem ajustes de dose.
Inibidores nucleosídeos/nucleotídeos da transcriptase reversa (INTIRs)		
Didanosina	Darunavir ↔ AUC ↔↓ Cmáx ↔ didanosina AUC ↑ didanosina Cmáx ↑	Darunavir/ritonavir e didanosina podem ser usados sem ajustes de dose. A didanosina deve ser tomada com o estômago vazio, portanto, deve ser tomada 1 hora antes ou 2 horas após darunavir/ritonavir administrado com alimentos.
Tenofovir disoproxil	Darunavir ↔ AUC ↑ Cmín ↑ Tenofovir tenofovir AUC ↑ Cmáx ↑ Tenofovir (↑tenofovir do efeito no transporte de MDR-1 nos túbulos renais).	O monitoramento da função renal pode ser indicado quando darunavir/ritonavir é administrado em combinação com tenofovir, particularmente em pacientes com doença renal ou sistémica subjacente, ou em pacientes tomando agentes nefrotóxicos.
Emtricitabina/ tenofovir alafenamida	Darunavir ↔ tenofovir alafenamida ↔ tenofovir ↑	A dose recomendada de emtricitabina/tenofovir alafenamida é de 200 mg/10 mg uma vez ao dia quando usado com darunavir/ritonavir.
Abacavir Emtricitabina Lamivudina Estavudina Zidovudina	Não estudado. Com base nas vias de eliminação do zidovudina, emtricitabina, estavudina e lamivudina, que são excretados principalmente por via renal, e do abacavir cujo metabolismo é mediado pelo CYP450, não são esperadas interações entre estes medicamentos e darunavir/ritonavir	Darunavir/ritonavir pode ser usado com esses INTRIs sem ajuste de dose.
Inibidores não nucleosídeos/nucleotídeos da transcriptase reversa (NNRTIs)		
Efavirenz	#Darunavir AUC ↓ Cmín ↓ Cmáx ↓ efavirenz AUC ↑ Cmín ↑ Cmáx ↑ (↑ efavirenz da indução do CYP3A)	A monitorização clínica da toxicidade do sistema nervoso central associada ao efavirenz pode ser indicada quando é por ser indicada quando darunavir/ritonavir é administrado em combinação com efavirenz. Efavirenz em combinação com darunavir/ritonavir comprimidos 400 mg/100 mg uma vez ao dia pode resultar em Cmín de darunavir abaixo do ideal. Se for utilizado efavirenz, deve ser utilizado em combinação com darunavir/ritonavir 600 mg/100 mg duas vezes por dia (vide seção 4.4).
Etravirina	Darunavir AUC ↓ Cmín ↓ ↔ Cmáx ↓ etravirina AUC ↓ Cmín ↓ Cmáx ↓	Darunavir e etravirina 200 mg duas vezes por dia podem ser usados sem ajustes de dose.
Nevirapina	#Darunavir ↔ as concentrações plasmáticas são consistentes com os dados históricos. nevirapina AUC ↑ Cmín ↑ Cmáx ↑ (↑ nevirapina da inibição do CYP3A)	Darunavir/ritonavir e nevirapina podem ser usados sem ajustes de dose.
Rilpivirina	Darunavir AUC ↔ Cmín ↓ Cmáx ↔ rilpivirina AUC ↑ Cmín ↑ Cmáx ↑	Darunavir/ritonavir e rilpivirina podem ser usados sem ajustes de dose.
Inibidores da protease (IPs) do HIV - sem dose basal de ritonavir adicional		
Atazanavir	#Darunavir AUC ↔ Cmín ↓ Cmáx ↔ atazanavir AUC ↔ Cmín ↑ Cmáx ↓	Darunavir/ritonavir e atazanavir podem ser usados sem ajustes de dose.
Indinavir	#Darunavir AUC ↑ Cmín ↑ Cmáx ↑ indinavir AUC ↑ Cmín ↑ Cmáx ↔	Quando usado em combinação com darunavir/ritonavir a dose de indinavir pode ser reduzida de 800 mg duas vezes ao dia para 600 mg duas vezes ao dia para controlar os efeitos colaterais.
Saquinavir	#Darunavir AUC ↓ Cmín ↓ Cmáx ↓ saquinavir AUC ↑ Cmín ↑ Cmáx ↓	Não é recomendado combinar darunavir/ritonavir com saquinavir.

Inibidor da protease (IP) do HIV - com dose basal de ritonavir adicional		
Lopinavir/ritonavir	Darunavir AUC ↓ Cmín ↓ Cmáx ↓ lopinavir AUC ↑ Cmín ↑ Cmáx ↓	Devido a uma diminuição de 40% na exposição de darunavir, não foram estabelecidas doses apropriadas da combinação. Assim, o uso concomitante de darunavir/ritonavir e o medicamento combinado lopinavir/ritonavir é contra-indicado (vide seção 4.3).
ANTAGONISTA DE CCR5		
Maraviroc	As concentrações de darunavir, ritonavir foram consistentes com os dados históricos maraviroc AUC ↑ Cmáx ↑	A dose de maraviroc deve ser de 150 mg duas vezes ao dia quando coadministrado com darunavir/ritonavir
ANTAGONISTA DO RECEPTOR ADRENÉRGICO α-1		
Alfuzosina	Espera-se que darunavir/ritonavir aumente as concentrações plasmáticas de alfuzosina (devido ao CYP3A).	A coadministração de darunavir/ritonavir e alfuzosina é contraindicada (vide seção 4.3).
ANTIANGINOSOS/ANTIARRÍTMICOS		
Disopiramida Flecainida Lidocaína (sistémica) Mexiletina Propafenona Quinidina Bepridil Dronedarona Ivabradina Quinidina Ranolazina	Não estudado. Espera-se que darunavir/ritonavir aumente as concentrações plasmáticas destes medicamentos (devido à inibição do CYP3A4 e CYP2D6)	Recomenda-se prescrição e monitorização da concentração terapêutica, se disponível, para estes medicamentos quando administrados concomitantemente com darunavir/ritonavir. A coadministração de darunavir/ritonavir com amiodarona, bepridil, dronedarona, ivabradina, quinidina ou ranolazina é contraindicada (vide seção 4.3).
Digoxina	Digoxina AUC ↑ Cmín ↑ (↑ digoxina da provável inibição de P-gp)	Como a digoxina tem um índice terapêutico estreito, recomenda-se que o menor dose possível de digoxina seja inicialmente prescrita quando tratando darunavir/ritonavir. A dose de digoxina deve ser cuidadosamente titulada para obter o efeito clínico desejado enquanto se avalia a estado clínico geral do paciente.
ANTIBIÓTICOS		
Claritromicina	#Darunavir AUC ↓ Cmín ↑ Cmáx ↓ claritromicina AUC ↑ Cmín ↑ Cmáx ↑ (↑ claritromicina da indução do CYP3A e possível inibição da P-gp)	Deve-se ter cuidado quando a claritromicina é combinada com darunavir/ritonavir. Para pacientes com disfunção renal, a informação do produto da claritromicina deve ser consultada para a dose recomendada.
Ritaminicos		
Rifampicina Rifapentina	Não estudado. A rifampicina e a rifampicina são fortes indutores do CYP3A e diminuem as concentrações de outros inibidores de proteases e podem diminuir as concentrações plasmáticas de darunavir e ritonavir (devido à indução do CYP3A).	A associação de rifampicina e darunavir/ritonavir é contra-indicada (vide seção 4.3).
Ritabutina	Darunavir AUC ↑ Cmín ↑ Cmáx ↑ ritabutina AUC ↑ Cmín ↑ Cmáx ↑ (A ritabutina é um indutor e substrato do CYP3A.)	É necessária uma redução da dose de ritabutina em 75% da dose usual de 300 mg/dia (para 150 mg em dias alternados) e aumento do monitoramento de eventos adversos relacionados à ritabutina em pacientes recebendo darunavir/ritonavir. Em caso de efeitos secundários, deve considerar-se um novo aumento da dose subsequente da ritabutina e a monitorização dos níveis sanguíneos. Com base no perfil de segurança de darunavir/ritonavir, o aumento da exposição ao darunavir na presença de ritabutina não justifica um ajuste de dose para darunavir/ritonavir.
ANTICOAGULANTES/INIBIDORES DA AGREGAÇÃO PLAQUETÁRIA		
Apixabana Edoxabana Rivaroxabana	Não estudados. A coadministração de darunavir/ritonavir com essas anticoagulantes pode aumentar as concentrações do anticoagulante e aumentar o risco de sangramento. (inibição de CYP3A e P-gp)	O uso de darunavir/ritonavir e dessas anticoagulantes não é recomendado.
Dabigatrana, Ticagrelor	A administração concomitante de darunavir/ritonavir com dabigatrana ou ticagrelor é contra-indicada (vide seção 4.3).	A coadministração de clopidogrel com darunavir/ritonavir não é recomendada.
Clopidogrel	Não estudado. O uso de clopidogrel com darunavir/ritonavir pode aumentar a exposição a dabigatrana ou ticagrelor e diminuir a exposição a um metabolito ativo de clopidogrel, o que pode reduzir a atividade antiplaquetária do clopidogrel.	O uso de anti-plaquetários não afetados pela inibição ou indução do CYP (*e.p.*, prasugrel).
ANTICONVULSIVANTES		
Fenobarbital Fenitoína	Não estudado. Prevê-se que o fenobarbital e a fenitoína diminuam as concentrações plasmáticas de darunavir (indução das enzimas CYP450).	Darunavir/ritonavir não deve ser usado em combinação com esses medicamentos.
Carbamazepina	Não é recomendado o ajuste de dose de darunavir/ritonavir. Se darunavir/ritonavir e carbamazepina precisarem ser coadministrados, os pacientes devem ser monitorados quanto a eventos adversos relacionados à carbamazepina. As concentrações de carbamazepina devem ser monitoradas, se possível, e sua dose titulada para a resposta adequada. A dose de carbamazepina pode precisar ser reduzida em 25-50% devido à presença de darunavir/ritonavir.	Recomenda-se o monitoramento clínico ao coadministrar darunavir/ritonavir e clonazepam.
Clonazepam	Não estudado A coadministração de darunavir/ritonavir com clonazepam pode aumentar as concentrações de clonazepam. (inibição do CYP3A)	Recomenda-se o monitoramento clínico.
ANTIDEPRESSIVOS		
Paroxetina	#Darunavir AUC ↔ Cmín ↔ Cmáx ↔ paroxetina AUC ↓ Cmín ↓ Cmáx ↓	Se forem coadministrados antidepressivos com darunavir/ritonavir, recomenda-se a titulação da dose de antidepressivo com base na avaliação da resposta antidepressiva. Além disso, os pacientes em dose estável de antidepressivos que iniciam o tratamento com darunavir/ritonavir devem ser monitorados quanto à resposta antidepressiva.
Sertralina	#Darunavir AUC ↔ Cmín ↓ Cmáx ↔ sertralina AUC ↓ Cmín ↓ Cmáx ↓	
Amitriptilina Desipramina Imipramina Nortriptilina Trazodona	A uso concomitante de darunavir/ritonavir e esses antidepressivos pode aumentar as concentrações do antidepressivo. (inibição de CYP2D6 e CYP3A) (inibição de CYP3A).	Recomenda-se o monitoramento clínico quando darunavir/ritonavir é coadministrado com esses antidepressivos e pode ser necessário ajuste da dose do antidepressivo.
ANTIEMÉTICOS		
Domperidona	Não estudada. A domperidona é principalmente metabolizada pelo CYP3A4.	A coadministração de domperidona com darunavir/ritonavir é contraindicada porque o risco de eventos adversos cardíacos ou do sistema nervoso central pode aumentar.
ANTIFÚNGICOS		
Voriconazol	Não estudado. Ritonavir pode diminuir as concentrações plasmáticas de voriconazol (indução de enzimas CYP450)	O voriconazol não deve ser combinado com darunavir/ritonavir a menos que uma avaliação dos benefícios e riscos justifique o uso voriconazol.
Cetoconazol	#Darunavir AUC ↑ Cmín ↑ Cmáx ↑ cetoconazol AUC ↑ Cmín ↑ Cmáx ↑	Justifica-se a precaução e recomenda-se o monitoramento clínico. Quando a coadministração for necessária, a dose diária de cetoconazol não deve exceder os 200 mg.
Fluconazol Itraconazol Posaconazol	Não estudado. Darunavir pode aumentar as concentrações plasmáticas destes medicamentos. Itraconazol ou posaconazol podem aumentar as concentrações de darunavir. (inibição de CYP3A e glicoproteína-P)	Justifica-se a precaução e recomenda-se o monitoramento clínico. Quando a coadministração for necessária, a dose diária de itraconazol não deve exceder os 200 mg.
Clotrimazol	Não estudado. O uso sistêmico concomitante de clotrimazol com darunavir/ritonavir pode aumentar as concentrações plasmáticas de darunavir e ritonavir. (inibição de CYP3A) O com base no modelo farmacocinético da coadministração.	É necessária precaução e recomenda-se o monitoramento clínico, quando é necessária a coadministração de clotrimazol. As frequências são definidas: muito comum (≥ 1/10); comum (≥ 1/100 a < 1/10); incomum (≥ 1/1.000 a < 1/100); rara (≥ 1/10.000 a < 1/1.000); muito rara (< 1/10.000) e desconhecida (não pode ser estimada a partir dos dados disponíveis).
MEDICAMENTOS ANTIGOTA		
Colchicina	Não estudado. O uso concomitante de colchicina e darunavir/ritonavir pode aumentar a exposição à colchicina (inibição do CYP3A e P-gp).	A redução da dose de colchicina ou a interrupção do tratamento com colchicina é recomendada em pacientes com função renal ou hepática normais se o tratamento com darunavir/ritonavir for necessário. Os doentes com comprometimento renal ou hepático não devem receber colchicina com darunavir/ritonavir (vide as seções 4.3 e 4.4).
ANTIMALÁRICOS		
Artemeter/lumefantrina	Darunavir AUC ↔ ↓ Cmín ↔ ↓ Cmáx ↔ ↓ artemeter e diidroartemisinina AUC ↓ Cmín ↔ ↓ Cmáx ↓ lumefantrina AUC ↑ Cmín ↑ Cmáx ↑	A associação de darunavir e artemeter/lumefantrina é contraindicada devido ao acordo da associação à lumefantrina.
Artemisinina	Pode ser necessário o ajuste de dose. Monitore o efeito clínico.	Pode ser necessário o ajuste de dose. Monitore o efeito clínico.
Halofantrina	Não estudado. A halofantrina é extensivamente metabolizada pelo CYP3A4. Espera-se que a inibição do metabolismo da halofantrina pelo ritonavir aumente a exposição à halofantrina, podendo potencialmente prolongar o intervalo QT...	A administração concomitante de darunavir potenciado e halofantrina está contra-indicada. A halofantrina tem um índice terapêutico estreito com um risco aumentado de prolongamento do intervalo QT em exposições mais elevadas.
Mefloquina	A coadministração pode aumentar a exposição à mefloquina.	Recomenda-se cautela e monitoramento rigoroso.
Proguanil	A coadministração pode diminuir a exposição ao proguanil.	A coadministração de atovaquona/proguanil deve ser evitada sempre que possível. Se for considerado clinicamente necessário, considere tomar atovaquona/proguanil com uma refeição rica em gordura para aumentar sua biodisponibilidade e monitorar a resposta do tratamento.
Quinina	A coadministração pode aumentar a exposição à quinina.	Além disso, recomenda-se cautela, pois a quinina aumenta o risco